



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

in certain significant germinal factors. Thus if *A* is sterile with *B* and *C*, *B* will be sterile with *C*. Shifting now<sup>8</sup> to the physiological technique, they discover that sterility is tied up with the rate of pollen-tube growth. In cases of selfing or cross-sterile pollinations, the pollen not only germinates successfully but develops a normal pollen tube. This tube grows through the style at a uniform rate, but fails to reach its goal before the flower decays. In cases of cross-fertility the pollen tube grows at a progressively increasing rate. The logical conclusion is that self-sterility is not due to the presence of inhibiting substances, but rather to the absence of accelerating substances (catalyzers produced by the pollen-tube nucleus in compatible crosses, and only local in their effect). At the wane of the flowering period self-sterility and cross-sterility may be replaced by "pseudo-fertility." This is explained by the breaking down of the stylar tissue, so that own pollen tubes grow at a uniformly greater (not accelerated) rate.—MERLE C. COULTER.

**Photometry.**—The probability that a solution of uranium acetate and oxalic acid may be used successfully as a chemical photometer in physiological researches involving the measurement and comparison of light intensities is indicated by some experiments by RIDGWAY.<sup>9</sup> The solution used consisted of 1 per cent uranium acetate and 5 per cent oxalic acid in aqueous solution mixed in the proportion of 1:4. In various tests designed to compare the oxalic-acid-uranium-salt photometer with the Callendar pyrheliometer, the chemical photometer gave results in general agreement with the pyrheliometer, even though the 2 instruments involve different portions of the solar spectrum. If the instrument and methods of using the solution can be reliably standardized, the inexpensiveness of the materials, ease of taking readings, accuracy of determinations, and its automatic integration for variable conditions of light during exposure will make it an excellent instrument for extending our knowledge of the influence of light as related to life processes in plants and animals.—C. A. SHULL.

**Nitrogen fixation.**—Another contribution from the Missouri Botanical Gardens on the subject of nitrogen fixation by lower organisms deals with the growth of *Azotobacter* in synthetic media. ALLEN<sup>10</sup> believes that most of the discrepancies in the results of previous investigators can be explained on the basis of the phosphorus requirements of the organism and the reaction of the medium. He proves this in a fairly satisfactory way by means of the following factors: (1) when  $\text{CaCO}_3$  is used to maintain a proper  $\text{P}_H$  in the media, the

<sup>8</sup> EAST, E. M., and PARK, J. B., Studies on self-sterile plants. II. Pollen-tube growth. *Genetics* 3:353-366. figs. 3. 1918.

<sup>9</sup> RIDGWAY, CHARLES S., A promising chemical photometer for plant physiological research. *Plant World* 21:234-240. 1918.

<sup>10</sup> ALLEN, E. R., Some conditions affecting the growth and activities of *Azotobacter chroococcum*. *Ann. Mo. Bot. Gard.* 6:1-44. 1919.